

## OS26lite Arthroscopy of the Shoulder

### Further information

You can get more information and share your experience at [www.aboutmyhealth.org](http://www.aboutmyhealth.org)

Tell us how useful you found this document at [www.patientfeedback.org](http://www.patientfeedback.org)



[www.rcseng.ac.uk](http://www.rcseng.ac.uk)



[www.bads.co.uk](http://www.bads.co.uk)



[www.rcsed.ac.uk](http://www.rcsed.ac.uk)



This document will give you information about an arthroscopy of the shoulder. If you have any questions, you should ask your GP or other relevant health professional.

### **What is an arthroscopy of the shoulder?**

An arthroscopy (also called 'keyhole' surgery) allows your surgeon to see inside your shoulder using a camera inserted through small cuts in the skin. They should be able to treat some problems using special surgical instruments.

### **What are the benefits of surgery?**

The main benefit of surgery is to confirm exactly what the problem is and, in many cases, to treat the problem at the same time.

### **Are there any alternatives to surgery?**

Problems inside a joint can often be diagnosed using tests such as CT and MRI scans. However, you may then need an arthroscopy to treat the problem.

### **What does the operation involve?**

The operation is usually performed under a general anaesthetic. The operation usually takes about forty minutes.

Your surgeon will usually make about two to four small cuts around the joint. They will place a small telescope through one of the cuts so they can examine the joint. They will place surgical instruments through the other cuts if they need to treat any problems with the joint (see figure 1).




Figure 1

Arthroscopy of the shoulder

## **What complications can happen?**

### **1 General complications**

- Pain
- Bleeding
- Infection of the surgical site (wound)
- Unsightly scarring

### **2 Specific complications**

- Bleeding into the joint
- Infection in the joint
- Severe pain, stiffness and loss of use of the arm and hand (complex regional pain syndrome)
- Damage to nerves
- Blood clot in the axillary vein near your armpit

### **How soon will I recover?**

You should be able to go home the same day.

Your physiotherapist may give you exercises and advice to help you to recover from the operation. It can take up to three months to get back to normal activities.

Regular exercise should help you to return to normal activities as soon as possible. Before you start exercising, you should ask a member of the healthcare team or your GP for advice.

Most people have a major improvement. However, it does take time for pain to lessen and movement to increase. Symptoms often come back with time.

### **Summary**

An arthroscopy of the shoulder allows your surgeon to diagnose and treat problems affecting the joint, without the need for a large cut in the skin. This may reduce the amount of pain you feel and speed up your recovery after surgery.

## **Acknowledgements**

Author: Prof. John Stanley MCh Orth FRCS (Ed.)  
FRCSE and Prof. Lennard Funk MSc FRCS (Tr. &  
Orth.)

Illustrations: Medical Illustration Copyright © 2011  
Nucleus Medical Art. All rights reserved.  
www.nucleusinc.com

**This document is intended for information  
purposes only and should not replace advice  
that your relevant health professional would  
give you.**